

Description: DNA molecules are built of dNTPs which are used in various PCR-based assays. The purity of dNTPs is highly important for the accuracy of assay results. The dNTPs synthesis itself doesn't except the presence of contaminants (such as NTPs, modified nucleotides, dNDPs, dNMPs, heavy/transition metals) in resulting solution, which can extremely affect the experiment by PCR inhibition.

The use of a highly purified dNTP preparation is particularly recommended for sensitive techniques such as long-range PCR, RT-PCR, multiplex, mutagenesis experiments and Real Time applications.

Content

Ref No.	110003	110004	color
dATP, 100 mM	200 μ L	1000 μ L	white
Datasheet	1	1	--

Applications: Many applications where high-quality reagents are required like reverse transcription (RT), polymerase chain reaction (PCR), RT-PCR, DNA labeling reactions, and sequencing/cycle sequencing analysis.

Concentration: In water of sodium salts of dATP 100 mM, pH 7.5, MW 634.2

Quality Control

- HPLC analysis (> 98 %);
- NMR analysis (inorganic phosphates)
- Exo-endonucleases contamination test
- UV-spectral analysis
- Spectrophotometry
- Production of 8 kb PCR fragment from genomic DNA with *Taq* DNA polymerase
- Production of 0.6 kb PCR fragment from genomic DNA with *Pfu* DNA polymerase

Storage condition: -20 °C

Note: The solution is ready-for-use and is optimized for PCR. Use 200 μ M of dATP solution in the PCR reaction volume.